



Anesthesiology Clin N Am
21 (2003) 835–857

ANESTHESIOLOGY
CLINICS OF
NORTH AMERICA

Cumulative Index 2003

Volume 21

March	ISSUES IN OBSTETRIC ANESTHESIA, pages 1–206
June	AMBULATORY ANESTHESIA, pages 207–451
September	ANESTHESIA AND THE CARDIAC PATIENT, pages 453–698
December	INTERVENTIONAL PAIN MANAGEMENT, pages 699–857

Note: Page numbers of article titles are in **boldface** type.

A

- Abdominal aortic aneurysms, endovascular exclusion of, in the cardiac catheterization laboratory, 648–650
- Abdominal delivery. See *Cesarean section*.
- Abruptio placenta, obstetric hemorrhage due to, 114–115
- Abscess, neuraxial, with neuraxial anesthesia in outpatient surgery, 296
- Acadesine, for neuroprotection in cardiac surgery, 489
ischemic preconditioning before CABG with, 478
- Accreditation, in office-based anesthesia, 431–433
- Acetaminophen, for pain management in ambulatory setting, 333–335
- Acetylcholinesterase inhibitors, intrathecal, 810
- Activated clotting time, monitoring of, for cardiopulmonary bypass, 512–514
- Acupuncture, for pain management in ambulatory setting, 342
potential for antiemetic use after ambulatory surgery, 356–357
- Acute normovolemic hemodilution, before delivery, in parturients at risk of hemorrhage, 132–133
- Adhesions, lysis of, for management of chronic back pain, **743–764**
caudal epidural compressive neuroplasty, 745–753
cervical epidural neuroplasty, 757–761
epiduroscopy, 755–757
history, 743–745
lumbar transforaminal approach, 753–755
thoracic epidural neuroplasty, 762–763
- Adrenocorticotrophic hormone, for postdural puncture headache, 76
- Air embolism, venous, during pregnancy, 170–173
- Air, intracardiac, detection of after valvular procedures with transesophageal echocardiography, 607–608
- Airway, difficult, obstetric emergencies due to, 119–120
- Airway management, in ambulatory anesthesia, Airway Management Device, 322–323
Cuffed Oropharyngeal Airway, 322
laryngeal mask airway for, 319–321
extended laparoscopy with, 320
in the prone position, 321
laparoscopic cholecystectomy with, 320–321
simple laparoscopy with, 319–320
Laryngeal Tube, 323
Streamlined Pharynx Airway Liner, 324
Airway Management Device (AMD), 322–323

- Allosteric modifiers, clinical potential during cardiac surgery, 555, 558-559
as transfusion alternative to red blood cells, 561
in stroke, 564
- Alpha-2 adrenergic agonists, intraspinal analgesia with, 807
- Ambulatory anesthesia, 207-444
controversial issues in, **313-327**
airway management, 319-324
depth-of-anesthesia monitoring, 313-319
development of, **207-228**
agents for, 210-213
financial and societal pressures, 221-226
cost reduction, 221-223
management challenges, 224-225
maximizing health care resources, 221
office-based surgery, 225-226
patient and surgeon satisfaction, 223-224
future technology, 219-220
information technology, 220
monitoring, 219-220
surgical, 219
history of emergence of, 208-210
patient selection, 217-219
processes, 215-217
techniques for, 213-215
inhalational techniques in, **263-272**
intravenous techniques in, **273-288**
office-based, overview of, **417-444**
patient outcomes, **403-415**
pediatric, current issues in, **305-311**
- Ambulatory surgery, challenging patients in setting of, **243-261**
elderly patients, 256-257
infants, 257-258
with aortic stenosis, 250-251
with end-stage renal disease, 255-256
with ischemic heart disease, 245-249
with obstructive sleep apnea, 252-255
effective analgesic modalities, **329-346**
postoperative nausea and vomiting, **347-365**
postoperative recovery and discharge, **367-386**
preoperative screening for, **229-242**
regional anesthesia for, **289-303**
scheduling of cases for, **387-401**
- Ambulatory surgical center, scheduling of cases in, **387-401**
- American Society of Anesthesiologists (ASA), Closed Claims database in obstetrics, lessons learned from, **183-198**
- Aminocaproic acid, epsilon-, use of during routine cardiopulmonary bypass, 531-532
- Amniotic fluid embolism, 173-179
- Analgesia, during labor, combined spinal-epidural, 5-10
continuous spinal, 4-5
epidural, patient-controlled, 10-12
patient-controlled, epidural, 10-12
inhalational, 2
intravenous, 1-2
effective modalities for ambulatory patients, **329-346**
clonidine, 338-339
dextromethorphan, 338
intraoperative, with intravenous anesthesia, 276-278
JCAHO standards for, 330-334
ketamine, 337-338
local anesthetics for, 339-342
local nerve blocks for, 341-342
NSAIDs and acetaminophen, 333-337
opioids, 337
others, 342
pain mechanisms and, 334-335
preemptive analgesia, 335
See also *Pain management and Analgesics*.
- Analgesics, spinal, **803-814**
experimental agents, 809-811
intrathecal acetylcholinesterase inhibitors, 810
nitric oxide synthase inhibitors, 810
sustained release delivery of local anesthetics, 810-811
tricyclic antidepressants, 810
intraspinal opioids, 804-806
fentanyl and sufentanil, 805
methadone and meperidine, 805
morphine, 805
side effects, 805-806
local anesthetics as adjuvant agents, 806
non-opioids, 806-809
alpha-2 adrenergic agents, 807-808
benzodiazepines, 809
calcium channel blockers, 808

- cells in matrix, 809
long-acting local anesthetics, 809
N-methyl-D-aspartate (NMDA) antagonists, 808-809
somatostatin analogs, 806
spinal infusion research, 811-812
tolerance and safety, 811
treatment criteria, 803-804
- Anaphylactoid syndrome of pregnancy, 173-179
- Anesthesia, ASA Closed Claims database on obstetric injuries due to, lessons learned from, **183-198**
for patients with severe pre-eclampsia, 156-160
general. See *General anesthesia*.
local. See *Local anesthesia*.
obstetric. See *Obstetric anesthesia*.
office-based, **417-444**
regional. See *Epidural anesthesia and Regional anesthesia*.
- Anesthesiologists, incomplete epidural block for cesarean section due to performance of, 46-47
performance of preoperative assessment for ambulatory surgery by, 236-237
- Anesthetic agents, for ambulatory surgery, developments in, 210-213
intravenous, 274-276
- Aneurysms, abdominal aortic, endovascular exclusion of, in the cardiac catheterization laboratory, 648-650
- Angina pectoris, spinal cord stimulation for, **795-802**
- Ankle block, in outpatient surgery, 293
- Antenatal assessment, reducing risk of obstetric hemorrhage by, 129-130
- Antenatal care, optimization of parturient to reduce risk of obstetric hemorrhage, 130-132
- Anti-inflammatory strategies, for cardiopulmonary bypass-induced inflammation, 458-462
- Anticoagulant medications, obstetric anesthesia for patients taking, **99-109**
- Anticoagulation, monitoring hemostasis and, for cardiopulmonary bypass, **511-526**
- Antidepressants, tricyclic, intraspinal analgesia with, 810
- Antiemetics, for use after ambulatory surgery, in clinical use, 349-355
with potential clinical use, 355-356
- Antifibrinolytics, in cardiopulmonary bypass (CPB), **527-551**
bleeding after CPB, 527-528
pathophysiology of excessive bleeding after CPB, 528-530
use during routine CPB, 530-541
use in elective deep hypothermic circulatory arrest, 541-543
- Antihistamines, for antiemetic use after ambulatory surgery, 352-353
- Antihypertensive agents, in pre-eclampsia, 154-156
- Antioxidants, for cardiopulmonary bypass-induced inflammation, 460
ischemic preconditioning before CABG with, 479
- Aortic regurgitation, intraoperative transesophageal echocardiography in, 600-602
- Aortic stenosis, ambulatory anesthesia in patients with, 250-251
intraoperative transesophageal echocardiography in, 599-600
- Aortic valve procedures, transesophageal echocardiography in, 596-603
after cardiopulmonary bypass, 602-603
before cardiopulmonary bypass, 596-602
- Apnea, obstructive sleep, ambulatory anesthesia in patients with, 252-255
- Aprotinin, for neuroprotection in cardiac surgery, 489
ischemic preconditioning before CABG with, 478-479
use of during routine cardiopulmonary bypass, 536-541
clinical safety, 540-541
efficacy, 537-540
mechanism of action, 536-537
trials of, 538-539
- Arachidonic acid metabolites, in systemic inflammatory response to cardiopulmonary bypass, 458
- Arrhythmias, in adults with congenital heart disease, 654, 656
perioperative, new treatments for, **569-586**

- basic science, 569-574
 - pediatric, 580-584
 - congenital heart disease and, 581-584
 - incidence and anesthetic factors, 581
 - supraventricular, 574-577
 - acute management, 574-576
 - chemical cardioversion of, 576-577
 - ventricular, 577-580
 - drug selection for acute management of, 579-580
 - nonsustained, 577-578
 - sustained, 578-579
 - Aspiration, pulmonary, of gastric contents during labor, decline in maternal mortality due to, 88-90
 - Assessment, preoperative. *See Preoperative assessment.*
 - Atony, uterine, obstetric hemorrhage due to, 112-114
 - Atrial septal defect, in adult patients with congenital heart disease, 654, 658
 - Atrioventricular canal, in adult patients with congenital heart disease, 655, 663
 - Audit, of cases of major obstetric hemorrhage, 139-140
 - Auditory evoked potential index, for depth-of-anesthesia monitoring in ambulatory anesthesia, 317-318
 - Autologous blood donation, preoperative, in parturients at risk of hemorrhage, 131-132
 - Automatic implantable cardioverter-defibrillators (AICDs), procedures with, in the cardiac catheterization laboratory, 647-648
 - Autonomic monitors, for depth-of-anesthesia monitoring in ambulatory anesthesia, 318
 - Axillary block, in outpatient surgery, 291-292
- B**
- Back pain, epidural analgesia and, **71-73**
 - chemical backache, 73
 - etiology, 71
 - incidence, 71-73
 - misconceptions about, 66
 - lysis of adhesions for management of, **743-764**
 - with neuraxial anesthesia in outpatient surgery, 295-296
 - Backache. *See Back pain.*
 - Benzamides, for antiemetic use after ambulatory surgery, 352
 - Benzodiazepines, intraspinal analgesia with, 809
 - Beta-adrenergic blocking agents, to control hypertension in pre-eclampsia, 154-155
 - Beta-blockers, for neuroprotection in cardiac surgery, 491
 - Bispectral index, for depth-of-anesthesia monitoring in ambulatory anesthesia, 314-317
 - and conscious sedation, 315
 - controversies in, 315-317
 - Bleeding, after cardiopulmonary bypass, 527-528
 - excess, pathophysiology of, 528-530
 - Bleeding time, point-of-care hemostasis monitors for after cardiopulmonary bypass, 520-523
 - Blocks, interventional pain management with, psychological aspects of, 769-774
 - Blood patch. *See Epidural blood patch.*
 - Blood salvage, intraoperative, for hemorrhage in obstetric patients, 134-135
 - Blood substitutes. *See Oxygen therapeutics.*
 - Blood transfusion. *See Transfusions, blood.*
 - Botulinum toxin therapy, in pain management, **713-729**
 - antibody formation, 716-717
 - clinical studies with, 717-723
 - dosing considerations, 723-725
 - history and early clinical development, 713-714
 - mechanism of action, 715
 - pharmacology, 714, 715-716
 - structure, 714-715
 - Breastfeeding, epidural analgesia and, misconceptions about, 64-66
 - Bupivacaine, compared with new local anesthetics for obstetric anesthesia, **19-38**
 - animal studies of, 20-25
 - chemistry of, 19, 26
 - clinical impact of toxicity of, 29-30
 - cost, 34
 - human studies, 28-29
 - motor block and, 30-31
 - relative potency, 31-34
 - toxicity of, 26-28
 - for spinal anesthesia in outpatient surgery, 297

Business aspects, of office-based anesthesia, 422-425

Butamben, intraspinal analgesia with, 809

Butyrophenones, for antiemetic use after ambulatory surgery, 351-352

C

C5 complement inhibitor, for neuroprotection in cardiac surgery, 492

Calcium channel blockers, intraspinal analgesia with, 808
to control hypertension in pre-eclampsia, 155

Calcium homeostasis, alterations in, in perioperative myocardial ischemia-reperfusion injury, 467-468

Cancer, effect of pain on survival in, 825-827
impact of psychotherapy on survival, 828

Cannabinoids, potential for antiemetic use after ambulatory surgery, 355

Cardiac arrest, maternal, obstetric emergencies due to, 123

Cardiac catheterization laboratory, anesthesia in, **639-651**
emergent care plan, 645-646
general *versus* regional techniques, 644-645
organization and arrangement of, 639-642
pharmacy availability and support services, 645
radiation safety, 643-644
radiocontrast media, 646-647
specific procedures, 647-650
 endovascular exclusion of abdominal aortic aneurysms, 648-650
 pacemaker and AICD changes and laser lead extractions, 647-648

Cardiac failure, left ventricular assist devices and, **625-638**

Cardiac surgery, 453-689
antifibrinolytics in cardiopulmonary bypass and elective deep hypothermic circulatory arrest, **527-551**
blood substitutes and oxygen therapeutics in, **553-568**
cardiac catheterization laboratory, anesthesia in, **639-651**
cardiac failure and left ventricular assist devices, **625-638**
for adults with congenital heart disease, **653-673**

gender and, **675-689**

monitoring anticoagulation and hemostasis in, **511-526**

neuroprotection in, **487-509**

off-pump, differences in anesthesia for, **613-623**

perioperative cardiac arrhythmias, new treatments for, **569-586**

perioperative myocardial ischemia reperfusion injury, **465-485**

systemic inflammatory response to cardiopulmonary bypass, **453-464**
transesophageal echocardiography in, **587-612**

Cardiopulmonary bypass (CPB), antifibrinolytics in, current status of, **527-551**
during routine CPB, 528-541
in elective deep hypothermic circulatory arrest, 541-543
intraoperative transesophageal echocardiography before and after, **587-612**
detection of intracardiac air with, 607-608
in aortic valve procedures, 596-603
in mitral valve procedures, 588-596
in tricuspid valve procedures, 604-607
potential role of oxygen therapeutics in, 562-563
systemic inflammatory response to, **453-464**
anti-inflammatory therapeutic strategies, 458-462
 mechanical, 461-462
 pharmacologic, 459-461
cellular response, 455-456
contact activation, 453-455
plasma inflammatory mediators, 457-458

Cardiovascular risk, ambulatory anesthesia in patients with, 248

Case scheduling, in an ambulatory surgical center, **387-401**

Catheter defects, incomplete epidural block for cesarean section due to, 44-46

Catheter malfunction, incomplete epidural block for cesarean section due to, 44-46

Catheter migration, incomplete epidural block for cesarean section due to, 44

Catheter misplacement, incomplete epidural block for cesarean section due to, 43-44

Catheterization, cardiac. *See* Cardiac catheterization laboratory.

- Cauda equina syndrome, after regional anesthesia, 82
- Caudal epidural decompressive neuroplasty, 745-753
See also *Adhesions, lysis of*.
- Caudal injections, supplemental, after incomplete epidural block for cesarean section, 49-50
- Cells in matrix, intraspinal analgesia with, 809
- Central nervous system, changes in, in pre-eclampsia, 151-153
- Cervical dystonia, with pain, botulinum toxin therapy for, 718
- Cervical epidural neuroplasty, 757-761
- Cesarean section, anesthesia for, in patients with severe pre-eclampsia, 158-160
 general, 159-160
 regional, 158-159
incomplete epidural block for, 39-57
 causes of
 anatomic considerations, 43
 patient-related factors, 46
 technical skills or
 performance factor, 46
 technique, methodology, and
 equipment-related
 factors, 43-46
 description of, 41-42
 management of, if diagnosed
 preoperatively, 47-50
 intraoperative, 50-52
- Chemical backache, after regional anesthesia, 73
- Chemokines, in systemic inflammatory response to cardiopulmonary bypass, 457
- Children, ambulatory anesthesia in. See *Pediatric ambulatory anesthesia*.
- Cholecystectomy, laparoscopic, laryngeal mask airway in, 320-321
- Cholinergic antagonists, for antiemetic use after ambulatory surgery, 350-351
- Chronic pain. See *Pain management, for chronic pain*.
- Clomethiazole, for neuroprotection in cardiac surgery, 492
- Clonidine, as adjunct for regional anesthesia in outpatient surgery, 298
 for pain management in ambulatory setting, 338-339
 intraspinal analgesia with, 807
- Closed Claims database, of obstetric anesthesia injury cases, lessons learned from, 183-198
- Clostridium botulinum*. See *Botulinum toxin therapy*.
- Coagulation abnormalities, in pre-eclampsia, 150-151
- Coagulation correction, before delivery, in parturients at risk of hemorrhage, 132
- Coagulation defects, obstetric anesthesia in patients with, 99-109
- Coagulation tests, for monitoring hemostasis after cardiopulmonary bypass, 518-519
- Coarctation of the aorta, in adult patients with congenital heart disease, 654, 659
- Coccydynia, sacral neuromodulation for, 787-792
 traditional medical and surgical therapies, 785-786
- Cognitive dysfunction, as opioid side effect, 705
- Cold therapy, continuous-flow, for pain management in ambulatory setting, 342
- Colorectal surgeons, use of office-based anesthesia by, 437
- Complement inhibitors, for cardiopulmonary bypass-induced inflammation, 460-461
 ischemic preconditioning before CABG with, 480-481
- Complement system, in systemic inflammatory response to cardiopulmonary bypass, 453-455
 role in perioperative myocardial ischemia-reperfusion injury, 471-474
- Complex regional pain syndrome, 731-742
 clinical features and epidemiology, 731-733
 clinical guidelines, 734-735
 interventional management, 735-736
 pathophysiology, 733-734
 rehabilitation, 736-739
 trophic changes and movement disorder, 733
- Complications, cardiac, in surgery in adult patients with congenital heart disease, 654, 656-658
- Congenital anomalies, noncardiac, in adults with congenital heart disease, 658

- Congenital heart disease, adult patients with, **653-673**
 anesthetic management of, 665-669
 cardiac complications, 654, 656-658
 specific cardiac lesions, 654-656, 658-665
 perioperative arrhythmias in patients with, 581-584
- Congestive heart failure, left ventricular assist devices and, **625-638**
- Conscious sedation, bispectral index monitoring and, 315
 intravenous ambulatory anesthesia with, 279-280
- Constipation, as opioid side effect, 703-705
- Continuous catheter techniques, for regional anesthesia in outpatient surgery, 294
- Continuous spinal analgesia, during labor, 4-5
- Continuous-flow cold therapy, for pain management in ambulatory setting, 342
- Convulsions, prevention of, in pre-eclampsia, 153
- Coronary artery bypass grafting (CABG), in women, gender differences, **675-689**
 off-pump, anesthetic considerations in, **613-623**
 differences in, 615-620
 techniques specific for, 620-622
 controlling heart rate, 620-621
 ischemic preconditioning, 621
 mechanical support of circulation, 621-622
 versus CABG with cardiopulmonary bypass, 622
 perioperative myocardial ischemia-reperfusion injury, **465-485**
 systemic inflammatory response to cardiopulmonary bypass in, **453-464**
- Coronary artery disease, ambulatory anesthesia in patients with, 245-249
 effect of pain on survival in, 823-825
- Corticosteroids, for cardiopulmonary bypass-induced inflammation, 459
 ischemic preconditioning before CABG with, 479
- Cosmetic surgery, use of office-based anesthesia in, 439-440
- Cost, of ropivacaine and levobupivacaine versus bupivacaine for obstetric anesthesia, 34
- Cost reduction, scheduling cases in an ambulatory surgery center, **387-401**
 with ambulatory anesthesia, 221-223
 with inhalational versus intravenous, 268-269
- Cuffed Oropharyngeal Airway (COPA), 322
- Cyclo-oxygenase-2 (COX-2) inhibitors, for pain management in ambulatory setting, 334, 336-337
- Cystitis, interstitial, sacral neuromodulation for, 787-792
 traditional medical and surgical therapies, 785-786
- Cytokines, in systemic inflammatory response to cardiopulmonary bypass, 457
-
- ## D
- Death, as patient outcome after ambulatory anesthesia, 404
- Deep hypothermic circulatory arrest, elective, use of antifibrinolytic agents in, 541-543
- Deep vein thrombosis, during pregnancy, 166-170
- Depth-of-anesthesia monitoring, in ambulatory anesthesia, **313-319**
 autonomic monitors, 318
 bispectral index, 314-317
 using auditory evoked potential index, 317-318
 using EEG signal, 317
- Dextran patch, epidural, for postdural puncture headache, 77
- Dextromethorphan, for neuroprotection in cardiac surgery, 490
 for pain management in ambulatory setting, 338
- Dialysis patients, ambulatory anesthesia in, 255-256
- Discharge, recovery and, after ambulatory surgery, **367-386**
 after regional anesthesia, 374-376
 factors delaying, 376-380
 other, 380
 postoperative nausea and vomiting, 378-380
 postoperative pain, 376-378
 voiding before, 373-374
 without tolerating oral fluids, 373
- Discography, interventional pain management with, psychological aspects of, 774-776
- Distal sciatic nerve block, in outpatient surgery, 293
- Dopamine antagonists, for antiemetic use after ambulatory surgery, 351-352

Drinking, during labor. See *Eating, during labor.*

Dystonia, with pain, botulinum toxin therapy
for, cervical, 718
focal, 718

E

Eating, during labor, NPO policies, scientific validation of, **87-98**

- decline in maternal mortality from pulmonary aspiration, 88-90
- gastrointestinal anatomy and physiology during pregnancy, 90-91
- indirect evidence on, 93-96
- metabolic advantages of, 91-92
- outcome of labor and, 92-93

Ebstein anomaly, in adult patients with congenital heart disease, 655, 662

Echocardiography, transesophageal, during valvular heart surgery, **587-612**

- aortic valve procedures, 596-603
- detection of intracardiac air, 607-608
- mitral valve procedures, 588-596
- tricuspid valve procedures, 604-607

Economic outcome, of ambulatory anesthesia, 411-412

Education, patient, prior to discharge after ambulatory surgery, 380-381

Efficiency, in the OR, scheduling cases in an ambulatory surgery center, **387-401**

Elderly patients, ambulatory anesthesia in, 256-257

Electroencephalogram (EEG) signal, for depth-of-anesthesia monitoring in ambulatory anesthesia, 317

Emboli reduction, for neuroprotection in cardiac surgery, 496-498

Embolism, during pregnancy, **165-182**

- amniotic fluid embolism, 173-179
- venous air embolism, 170-173
- venous thromboembolism, 166-170

Emergencies, obstetric, **111-125**

- fetal compromise, 117-119
- hemorrhage, 111-117

- abruptio placenta, 114-115
- genital tract lacerations, 111-112
- placenta accreta, 116
- placenta previa, 115-116

- retained placenta, 115
- uterine atony, 112-114
- uterine inversion, 117
- uterine rupture, 116-117
- special considerations, 119-123
 - difficult airway, 119-120
 - maternal cardiac arrest, 123
 - morbid obesity, 121-123
 - pregnancy-induced hypertension, 120-121

Emergent care plan, for the cardiac catheterization laboratory, 645-646

Emesis, mechanism of, 347-349

End-stage renal disease, ambulatory anesthesia in patients with, 255-256

Endothelial activation, in systemic inflammatory response to cardiopulmonary bypass, 456

Endothelial cell-leukocyte interactions, in perioperative myocardial ischemia-reperfusion injury, 468-471

Endothelin-1, in systemic inflammatory response to cardiopulmonary bypass, 458

Endotoxemia, in systemic inflammatory response to cardiopulmonary bypass, 458

Endovascular exclusion, of abdominal aortic aneurysms, in the cardiac catheterization laboratory, 648-650

Epididymo-orchalgia, sacral neuromodulation for, 787-792

- traditional medical and surgical therapies, 785-786

Epidural analgesia, during labor, combined spinal and, 5-10

- patient-controlled, 10-12
- for labor and vaginal delivery in patients with severe pre-eclampsia, 157-158
- misconceptions about, **59-70**
 - back pain and, 66, 71-73
 - breastfeeding and, 64-66
 - consequences of withholding, 66-68
 - pre-eclampsia and, 63-64
 - progress of labor, 60-63

Epidural anesthesia, in outpatient surgery, 298-299

- incomplete block for cesarean section, **39-57**
 - causes of
 - anatomic considerations, 43
 - patient-related factors, 46
 - technical skills or performance factor, 46

- technique, methodology, and equipment-related factors, 43-46
- description of, 41-42
- management of, if diagnosed preoperatively, 47-50
- intraoperative, 50-52
- Epidural blood patch, for postdural puncture headache, 77
 - dextran, 77
 - prophylactic, 77
 - treatment with, 77-79
 - in HIV-positive patients, 79
- Epidural decompressive neuroplasty. See also *Adhesions, lysis of*
 - caudal, 745-753
 - cervical, 757-762
 - lumbar transforaminal approach, 753-755
 - thoracic, 762-764
- Epiduroscopy, 755-757
- Epinephrine, as adjunct for regional anesthesia in outpatient surgery, 298
- Epsilon-aminocaproic acid, use of during routine cardiopulmonary bypass, 531-532
- Erythropoietin, recombinant human, antenatal, in parturients at risk of hemorrhage, 130-131
- Esmolol, to control hypertension in pre-eclampsia, 155
- Estrogen replacement therapy, and gender differences in cardiac surgery, 679-682
- Etomidate, intravenous ambulatory anesthesia with, 275

F

- Factor XII, in systemic inflammatory response to cardiopulmonary bypass, 455
- Failed epidural. See *Epidural anesthesia, incomplete block*.
- Femoral block, in outpatient surgery, 293
- Fentanyl, as adjunct for regional anesthesia in outpatient surgery, 298
 - for intraoperative analgesia in intravenous ambulatory anesthesia, 276-277
 - intraspinal, 805
 - See also *Opioids*.
- Fetal compromise, obstetric emergencies involving, 117-119
 - choice of anesthesia, 118-119
 - definition, 117-118
 - in utero resuscitation, 118
 - neonatal resuscitation, 119
- Fibrin glue injection, epidural, for postdural puncture headache, 77
- Fibrinolysis, pathophysiology of excessive bleeding after cardiopulmonary bypass, 529
- Financial issues, in ambulatory anesthesia, cost reduction, 221-223
 - economic outcomes, 411-412
 - maximizing health care resources, 221
- Fluid management, to control hypertension in pre-eclampsia, 155-156
- Focal dystonia, with pain, botulinum toxin therapy for, 718
- Functional health status, as patient outcome after ambulatory anesthesia, 409-410

G

- Ganglion blocking agents, to control hypertension in pre-eclampsia, 154
- Gastric contents, pulmonary aspiration of during labor, decline in maternal mortality due to, 88-90
- Gastroenterologists, use of office-based anesthesia by, 437
- Gastrointestinal anatomy, during pregnancy, 90-91
- Gender differences, cardiac surgery in women, 675-689
 - operative factors and mortality, 682-683
 - pharmacology and estrogen replacement therapy, 679-682
 - postoperative long-term follow-up, 683-684
 - preoperative risk profile and mortality, 675-678
 - role of bias, 678-679
- General anesthesia, after incomplete epidural block for cesarean section, 50
 - and development of ambulatory anesthesia, 213
 - for cesarean delivery in patients with severe pre-eclampsia, 159-160
 - increased use of, due to withholding of epidural analgesia during labor, 67

- versus regional techniques, in cardiac catheterization laboratory, 644-645
- Genital tract lacerations, obstetric hemorrhage due to, 112
- Ginger, potential for antiemetic use after ambulatory surgery, 357
- GM1 ganglioside, for neuroprotection in cardiac surgery, 490
- Great arteries, transposition of, in adult patients with congenital heart disease, 660-661, 665
congenitally corrected, 655, 661
- Gynecology, use of office-based anesthesia in, 437
- H**
- Headache, postdural puncture, after regional analgesia during labor, 73-79
postprocedural, with spinal anesthesia in outpatient surgery, 295
- Heart disease, ambulatory anesthesia in patients with, aortic stenosis, 250-251
ischemic, 245-249
- Hematologic disease, in adults with congenital heart disease, 658
- Hematoma, neuraxial, with neuraxial anesthesia in outpatient surgery, 296
- Hemodilution, acute normovolemic, before delivery, in parturients at risk of hemorrhage, 132-133
- Hemodynamic profile, in pre-eclampsia, 146-153
- Hemofiltration, for cardiopulmonary bypass-induced inflammation, 461
- Hemoglobin-based oxygen carriers, clinical potential during cardiac surgery, 554, 556-557
as transfusion alternative to red blood cells, 559-560
in cardiopulmonary bypass, 562-563
- Hemorrhage, obstetric, 111-117
abruptio placenta, 114-115
genital tract lacerations, 111-112
placenta accreta, 116
placenta previa, 115-116
retained placenta, 115
steps to limit and treat, 127-144
antenatal assessment, 129-130
antenatal optimization of parturient, 130-132
at time of delivery, 133-136
audit and review of cases, 139-140
further treatment after stabilization, 138
future alternatives to blood transfusion, 138-139
immediately before delivery, 132-133
intraoperative management, 136-138
risks of blood transfusion, 128
uterine atony, 112-114
uterine inversion, 117
uterine rupture, 116-117
- Hemostasis, intravascular, intraoperative maintenance of, in hemorrhage in obstetric patients, 136-137
monitoring of, for cardiopulmonary bypass, 511-526
bleeding time and point-of-care monitors, 520-523
coagulation tests, 518-519
platelet function, 519-520
pathophysiology of excessive bleeding after cardiopulmonary bypass, 528-529
- Heparin coated circuits, to reduce cardiopulmonary bypass-induced inflammation, 462
- Heparin concentration, monitoring of, for cardiopulmonary bypass, 515-516
- Heparin effect, monitoring of, for cardiopulmonary bypass, 511-512
- Heparin management test, monitoring of, for cardiopulmonary bypass, 516-517
- Heparin neutralization, monitoring of, for cardiopulmonary bypass, 517-518
- Heparin resistance, monitoring of, for cardiopulmonary bypass, 514
- Heparin sensitivity measurement, monitoring of, for cardiopulmonary bypass, 514-515
- Heparin, low-molecular-weight, for venous thromboembolism during pregnancy, 169-170
obstetric anesthesia for patients on, 99-109
- Hepatic dysfunction, in pre-eclampsia, 149
- High dose thrombin time, monitoring of, for cardiopulmonary bypass, 516
- History, of ambulatory anesthesia, 207-226
- Hormone replacement therapy, and gender differences in cardiac surgery, 679-682

- Human immunodeficiency virus (HIV), epidural blood patch for postdural puncture headache in patients with, 79
- Hydralazine, to control hypertension in pre-eclampsia, 154
- Hyperbaric oxygen, for venous air embolism in pregnancy, 172-173
- Hypertension, pre-eclampsia, fluids, drugs and anesthetic management, **145-163**
 - pregnancy-induced, obstetric emergencies due to, 120-121
 - pulmonary, in adults with congenital heart disease, 656-657
- Hypnotic drugs, for intravenous ambulatory anesthesia, 274-276
- Hypogonadism, as opioid side effect, 706-709
- Hypothermia, elective deep hypothermic circulatory arrest, use of antifibrinolytics with, 541-543
 - for cardiopulmonary bypass-induced inflammation, 461
 - for neuroprotection in cardiac surgery, 493-496

I

- Ibuprofen, for pain management in ambulatory setting, 334, 335
- Imaging, intraoperative transesophageal echocardiography, **587-612**
 - myocardial contrast, potential role of oxygen therapeutics in, 561-562
- Implantable therapies. See also *Spinal cord stimulation*., interventional pain management with, psychological aspects of, 776-777
- In utero resuscitation, in obstetric emergencies involving fetal compromise, 118-119
- Incomplete epidural block. See *Epidural anesthesia*.
- Indomethacin, for pain management in ambulatory setting, 334, 335
- Induction of anesthesia, in ambulatory anesthesia, inhalational techniques for, 263-265
- Information technology, in ambulatory anesthesia, future developments in, 220
- Intraclavicular block, in outpatient surgery, 291
- Infusions, spinal, with implanted pump, research on, 811-812
- Inhalational analgesia, patient-controlled, during labor, 2
- Inhalational techniques, in ambulatory anesthesia, **263-272**
 - costs of, 268-269
 - induction, 263-265
 - laryngeal mask airway and, 268
 - maintenance, 265-266
 - nitrous oxide and, 267
 - postoperative nausea and vomiting with, 266-267
- Injuries, due to obstetric anesthesia, ASA Closed Claims database on, lessons learned from, **183-198**
- Interscalene anesthesia, in outpatient surgery, 290-291
- Interstitial cystitis, sacral neuromodulation for, 787-792
 - traditional medical and surgical therapies, 785-786
- Interventional pain management, psychological aspects of, **765-781**
 - blocks, 769-774
 - differential spinal blocks, 774
 - implantable therapies, 776-777
 - in discography, 774-776
 See also *Pain management*.
- Intraaortic balloon pump, in congestive heart failure, 627
- Intracardiac air, detection of after valvular procedures with transesophageal echocardiography, 607-608
- Intraoperative blood salvage, for hemorrhage in obstetric patients, 134-135
- Intraoperative management, of inadequate epidural block for cesarean section, 50-52
- Intraoperative transesophageal echocardiography. See *Transesophageal echocardiography*.
- Intrathecal drug delivery, of opioids, impact on survival, 829
 - See also *Spinal analgesia*.
- Intravenous analgesia, patient-controlled, during labor, 1-2
- Intravenous techniques, in ambulatory anesthesia, **273-288**
 - advantages of, 281-283
 - conscious sedation, 280-281
 - hypnotic drugs for, 274-276
 - intraoperative analgesia, 276-278
 - limits of, 283-284
 - monitored anesthesia care (MAC), 279-280
 - patient-controlled sedation, 281

- target-controlled infusion (TCI), 279
 - total intravenous anesthesia (TIVA), 278-279
 - Ion-channel mechanisms, in preoperative arrhythmias, 569-572
 - Iron supplementation, antenatal, in parturients at risk of hemorrhage, 130-131
 - Ischemia-reperfusion injury, perioperative myocardial, **465-485**
 - pathophysiology of, 466-474
 - treatment of, 475-481
 - Ischemic heart disease, ambulatory anesthesia in patients with, 245-249
 - Ischemic preconditioning, in off-pump coronary artery bypass, 621
 - to diminish perioperative myocardial ischemia-reperfusion injury, 477-481
-
- J**
- Joint Commission on Accreditation of Healthcare Organizations (JCAHO), pain standards for hospitals and ambulatory surgery centers, 330-331
-
- K**
- Kallikrein, in systemic inflammatory response to cardiopulmonary bypass, 455
 - Ketamine, for intraoperative analgesia in intra-venous ambulatory anesthesia, 277-278
 - for pain management in ambulatory setting, 337-338
 - Ketorolac, for pain management in ambulatory setting, 334, 335-336
-
- L**
- Labetalol, to control hypertension in pre-eclampsia, 154-155
 - Labor, newer techniques of analgesia during, **1-17**
 - NPO during, guidelines for, **87-98**
 - See also *Obstetric anesthesia*.
 - Labor costs, scheduling cases in an ambulatory surgery center, **387-401**
 - Laboratory testing, preoperative, for ambulatory surgery, 237-241
 - Lacerations, genital tract, obstetric hemorrhage due to, 112
 - Laparoscopy, laryngeal mask airway in, 319-321
 - cholecystectomy, 320-321
 - extended, 320
 - in the prone position, 322
 - simple, 319-320
 - Laryngeal mask airway, in ambulatory anesthesia, 268, 319-321
 - extended laparoscopy with, 320
 - in the prone position, 321
 - laparoscopic cholecystectomy with, 320-321
 - simple laparoscopy with, 319-320
 - Laryngeal Tube, 323
 - Laser extraction, of AICD leads, in the cardiac catheterization laboratory, 648
 - Left ventricular assist devices, **625-638**
 - anesthetic management for insertion of, 633-637
 - historical evolution and indications, 626-627
 - physiology and preoperative considerations, 630-632
 - surgical approach, 633
 - types of, 627-630
 - counterpulsation, 627
 - extracorporeal kinetic displacement, 627-628
 - nonpulsatile pumps, 629-630
 - pulsatile pumps, 628-629
 - Left ventricular outflow obstruction, intraoperative transesophageal echocardiography in, 595-596
 - Legislative issues, in office-based anesthesia, 428-430
 - Leukocyte activation, in systemic inflammatory response to cardiopulmonary bypass, 455-456
 - Leukocyte filters, for cardiopulmonary bypass-induced inflammation, 461
 - Leukocyte-endothelial cell interactions, in perioperative myocardial ischemia-reperfusion injury, 468-471
 - Levobupivacaine, compared with bupivacaine for obstetric analgesia, **19-38**
 - Lexiphan, for neuroprotection in cardiac surgery, 492
 - Liability, medico-legal, ASA Closed Claims database on obstetric anesthetic injuries, lessons learned from, **183-198**
 - Lidocaine, for neuroprotection in cardiac surgery, 491
 - for spinal anesthesia in outpatient surgery, 296-297

- Local anesthetics, for pain management in ambulatory setting, 339-341
 continuous infusion of, 341
 infiltrate, 340
 intraarticular, 341
 intraperitoneal, 340
intraspinal analgesia with
 long-acting, 809
 sustained-release delivery of, 810-811
new, for labor and delivery, **19-38**
 chemistry of, 19, 26
 compared with bupivacaine, animal studies of, 20-25
 cost of, 34
 human studies, 28-29
 reduced motor block with, 30-31
 relative potency of, 31-34
 toxicity, 26-28
- Local infiltration anesthesia, to supplement incomplete epidural block for cesarean section, 50
- Local nerve blocks, for pain management in ambulatory setting, 341-342
- Low back pain. See *Back pain*.
- Low-molecular-weight heparin, for venous thromboembolism during pregnancy, 169-170
 obstetric anesthesia for patients on, **99-109**
- Lower extremity blocks, in outpatient surgery, 292-293
- Lumbar sympathetic block, during labor, 3
- Lysine analogues, use of during routine cardiopulmonary bypass, 531
- Lysis, of adhesions. See *Adhesions*, lysis of.
-
- M**
- Magnesium sulfate, for prevention of convulsions in pre-eclampsia, 153
- Magnetic resonance imaging (MRI), and neuroprotection in cardiac surgery, 500-501
- Maintenance of anesthesia, in ambulatory anesthesia, inhalational techniques for, 265-266
- Malpractice claims, ASA Closed Claims database on claims involving obstetric anesthesia, **183-198**
- Management issues, in ambulatory surgery, 224-225
 scheduling of cases, **387-401**
- Maternal cardiac arrest, obstetric emergencies due to, 123
- Maternal death, due to pulmonary aspiration of gastric contents during labor, decline in, 88-90
- Maternal obesity, obstetric emergencies due to, 121-123
- Mechanical support, of circulation, in off-pump coronary artery bypass, 621-622
- Meperidine, intraspinal, 805
 See also *Opioids*.
- Methadone, intraspinal, 805
 See also *Opioids*.
- Methohexital, intravenous ambulatory anesthesia with, 274
- Midazolam, intravenous ambulatory anesthesia with, 275
- Mitral regurgitation, intraoperative transesophageal echocardiography in, 591-593
- Mitral stenosis, intraoperative transesophageal echocardiography in, 589-591
- Mitral valve procedures, transesophageal echocardiography in, 588-596
 after cardiopulmonary bypass, 593-596
 before cardiopulmonary bypass, 588-593
- Monitored anesthesia care (MAC), development of, for ambulatory surgery, 214-215
 in cardiac catheterization laboratory, 644-645
 intravenous ambulatory anesthesia with, 279-280
- Monitoring, in ambulatory anesthesia, future developments in, 219-220
- Mood, interaction of pain and, 827
- Morbid obesity, maternal, obstetric emergencies due to, 121-123
- Morbidity, non-anesthetic-related, as patient outcome after ambulatory anesthesia, 404-405
- Morphine, intraspinal, 805
 See also *Opioids*.
- Mortality, in cardiac surgery, gender differences in, 675-678
 operative factors and, 682-683
 maternal, due to obstetric anesthesia, ASA Closed Claims database on, 184-187

due to pulmonary aspiration of gastric contents during labor, decline in, 88-90

Motor block, in obstetric anesthesia, with bupivacaine *versus* bupivacaine, 30-31

Myocardial contrast imaging, potential role of oxygen therapeutics in, 561-562

Myocardial ischemia-reperfusion injury, perioperative, **465-485**
pathophysiology of, 466-474
treatment of, 475-481

Myoclonus, as opioid side effect, 706

Myofascial pain syndrome, botulinum toxin therapy for, 722-723

N

N-Methyl-D-aspartate antagonists, intraspinal analgesia with, 808-809

Na⁺/H⁺ exchange (NHE) inhibitors, ischemic preconditioning before CABG with, 481

Nausea and vomiting, as opioid side effect, 703
postoperative, management of, after ambulatory surgery, **347-365**

Needle-induced trauma, nerve injury due to, after regional anesthesia, 80-81

Neonatal resuscitation, in obstetric emergencies involving fetal compromise, 119

Nerve blocks, local, for pain management in ambulatory setting, 341-342

Nerve injury, from needle, after regional anesthesia, 80-81

Neuraxial analgesia., in patients with thrombocytopenia, **99-109**
misconceptions about, **59-70**
See also *Epidural analgesia*.

Neuraxial techniques, for regional anesthesia in outpatient surgery, 294-295

Neurokinin-1 antagonists, potential for antiemetic use after ambulatory surgery, 355-356

Neurologic complications, after regional anesthesia during labor, **79-82**
cauda equina syndrome, 82
etiology of, 79-80
nerve injury from needle, 80-81
transient neurologic syndrome, 81-82

Neurologic disease, in adults with congenital heart disease, 658

Neurologic symptoms, with neuraxial anesthesia in outpatient surgery, 295-296

Neuromodulation, sacral, for chronic pain conditions, **783-793**

Neuroprotection, in cardiac surgery, **487-509**
delayed postoperative stroke, 499-500
emboli reduction, 496-498
future of, 501-502
hypothermia, 493-496
magnetic resonance imaging, 500-501
off-pump cardiac surgery, 498-499
pharmacologic, 487-493

Neurostimulators. See *Spinal cord stimulation*.

Nicardipine, to control hypertension in pre-eclampsia, 155

Nifedipine, to control hypertension in pre-eclampsia, 155

Nil per mouth (NPO), during labor, guidelines for, **87-98**
decline in maternal mortality from pulmonary aspiration of gastric contents, 88-90
gastrointestinal anatomy and physiology during pregnancy, 90-91
indirect evidence on, 93-96
metabolic advantages of eating and drinking during labor, 91-92
outcome of labor, 92-93

Nimodipine, for neuroprotection in cardiac surgery, 490

Nitric oxide, in systemic inflammatory response to cardiopulmonary bypass, 458

Nitric oxide donors, for cardiopulmonary bypass-induced inflammation, 460

Nitric oxide synthase inhibitors, intraspinal analgesia with, 810

Nitroglycerin, to control hypertension in pre-eclampsia, 154

Nitrous oxide, and inhalation techniques in ambulatory anesthesia, 267

NMDA antagonists, intraspinal analgesia with, 808-809

Nonsteroidal anti-inflammatory drugs (NSAIDs), for pain management in ambulatory setting, 333-337

Normovolemic hemodilution, acute, before delivery, in parturients at risk of hemorrhage, 132-133

NPO. See *Nil per mouth*.

O

- Obesity, incomplete epidural block for cesarean section due to, 46
maternal morbid, obstetric emergencies due to, 121-123
- Obstetric anesthesia, 1-198
anticoagulant medications and, **99-109**
low-molecular-weight heparin, 105-107
practical recommendations, 104-105
ASA closed claims, lessons learned from, **183-198**
back pain after regional, **71-73**
embolism during pregnancy, **165-182**
amniotic fluid embolism, 173-179
venous air embolism, 170-173
venous thromboembolism, 166-170
headache after regional, **73-79**
hemorrhage, steps to limit and treat, **127-144**
antenatal assessment, 129-130
antenatal optimization of parturient, 130-132
at time of delivery, 133-136
audit and review of cases, 139-140
further treatment after stabilization, 138
future alternatives to blood transfusion, 138-139
immediately before delivery, 132-133
intraoperative management, 136-138
risks of blood transfusion, 128
in patients with thrombocytopenia, **99-109**
low-molecular-weight heparin and, 105-107
practical recommendations, 104-105
in pre-eclampsia, **145-163**
anesthetic management, 156-160
epidural analgesia and, misconceptions about, 63-64
hemodynamic profile in, 146-153
treatment, 153-156
incomplete epidural block for cesarean section, **39-57**
causes of
anatomic considerations, 43
patient-related factors, 46
technical skills or performance factor, 46
technique, methodology, and equipment-related factors, 43-46
description of, 41-42
management of, if diagnosed preoperatively, 47-50
intraoperative, 50-52
neuraxial analgesia, misconceptions about, **59-70**
back pain and, 66, **71-73**
breastfeeding and, 64-66
consequences of withholding, 66-68
pre-eclampsia and, 63-64
progress of labor, 60-63
neurologic complications after regional, **79-82**
new local anesthetics, **19-38**
chemistry of, 19, 26
compared with bupivacaine, animal studies of, 20-25
cost of, 34
human studies, 28-29
reduced motor block with, 30-31
relative potency of, 31-34
toxicity, 26-28
newer techniques of labor analgesia, **1-17**
patient-controlled analgesia
epidural, 10-12
inhalational, 2-3
intravenous, 1-2
regional techniques, 3-13
combined spinal-epidural analgesia, 5-10
continuous spinal analgesia, 4-5
epidural analgesia during labor, 10
lumbar sympathetic block, 3
paracervical block, 3-4
NPO during labor, guidelines for, **87-98**
decline in maternal mortality from pulmonary aspiration of gastric contents, 88-90
gastrointestinal anatomy and physiology during pregnancy, 90-91
indirect evidence on, 93-96
metabolic advantages of eating and drinking during labor, 91-92
outcome of labor, 92-93
obstetric emergencies and, **111-125**
fetal compromise, 117-119
hemorrhage, 111-117
abruptio placenta, 114-115
genital tract lacerations, 111-112
placenta accreta, 116
placenta previa, 115-116
retained placenta, 115
uterine atony, 112-114

- uterine inversion, 117
 - uterine rupture, 116-117
 - special considerations, 119-123
 - difficult airway, 119-120
 - maternal cardiac arrest, 123
 - morbid obesity, 121-123
 - pregnancy-induced
 - hypertension, 120-121
 - regional, backache, headache, and neurologic deficit after, **71-86**
 - Obstetrics, use of office-based anesthesia in, 437
 - Obstructive sleep apnea, ambulatory anesthesia in patients with, 252-255
 - Off-pump coronary artery bypass, anesthetic considerations, **613-623**
 - differences in, 615-620
 - neuroprotection with, 498-499
 - techniques specific for, 620-622
 - controlling heart rate, 620-621
 - ischemic preconditioning, 621
 - mechanical support of circulation, 621-622
 - versus conventional bypass graft with cardiopulmonary bypass, 622
 - Office-based anesthesia, overview of, 225-226, **417-444**
 - accreditation, 431-433
 - business aspects, 422-425
 - clinical essentials, 433-436
 - legislative progress, 428-430
 - professional recognition, 418-419
 - providers, 430-431
 - recovery, 436-437
 - safety issues, 426-428
 - specialties using, 437-440
 - specific attributes, 419-422
 - Oliguria, in pre-eclampsia, 149
 - Operating room (OR) management, scheduling of cases in an ambulatory center, **387-401**
 - Ophthalmology, use of office-based anesthesia in, 437-438
 - Opioids, for chronic pain, **699-712**
 - effectiveness of, 699-702
 - drug-centered characteristics, 701-702
 - pain-centered characteristics, 700-701
 - side effects of, 702-709
 - cognitive dysfunction, 705
 - constipation, 703-705
 - general principles, 702-709
 - hypogonadism, 706-709
 - myoclonus, 706
 - nausea or vomiting, 703
 - organ toxicity, 706
 - pruritus, 706
 - sedation, 705
 - source of, 702
 - strategies, 703
 - for pain management in ambulatory setting, 337
 - intraspinal, 804-806
 - fentanyl and sufentanil, 805
 - methadone and meperidine, 805
 - morphine, 805
 - side effects, 805-806
 - Organ toxicity, as opioid side effect, 706
 - Orthopedic surgery, use of office-based anesthesia in, 440
 - Otolaryngology, use of office-based anesthesia in, 438
 - Outcomes assessment, of therapeutic and cost benefit success rates of spinal cord stimulation, **815-821**
 - See also *Patient outcomes*.
 - Outpatient surgery. See *Ambulatory surgery*.
 - Oxygen, hyperbaric, for venous air embolism in pregnancy, 172-173
 - potential for antiemetic use after ambulatory surgery, 356
 - Oxygen therapeutic agents, clinical potential during cardiac surgery, **553-568**
 - allosteric modifiers, 558-559
 - hemoglobin-based oxygen carriers, 556-557
 - oxygen delivery to tissues, 553-556
 - perfluorochemical solutions, 557-558
 - potential applications, 559-564
 - as protection against ischemia and reperfusion injury, 563-564
 - in cardiopulmonary bypass, 562-563
 - in myocardial contrast imaging, 561-562
 - in stroke, 564
 - transfusion alternative to red blood cells, 559-561
 - Oxytocics, intraoperative, for hemorrhage in obstetric patients, 136
-
- P**
- Pacemaker changes, in the cardiac catheterization laboratory, 647-648

- Pain, components of, 766-767
 definitions of, 766
 effect of on survival, **823-831**
 impact of intrathecal opioid delivery, 829
 impact of pain management therapies-829, 828
 impact of psychotherapy on, 827-828
 in cancer, 825-827
 in coronary artery disease, 823-825
 in sickle cell disease, 825
 interaction of pain and mood, 827
 types of, 767-769
- Pain management, effective modalities for ambulatory patients, **329-346**
 clonidine, 338-339
 dextromethorphan, 338
 intraoperative, with intravenous anesthesia, 276-278
 JCAHO standards for, 330-334
 ketamine, 337-338
 local anesthetics for, 339-342
 local nerve blocks for, 341-342
 NSAIDs and acetaminophen, 333-337
 opioids, 337
 others, 342
 pain mechanisms and, 334-335
 preemptive analgesia, 335
 for chronic pain, 699-831
 botulinum toxin therapy, **713-729**
 complex regional pain syndrome, **731-742**
 effect of pain on survival, **823-831**
 lysis of adhesions, **743-764**
 opioid effectiveness and side effects, **699-712**
 psychological aspects of interventional therapy, **765-781**
 sacral neuromodulation, **783-793**
 spinal analgesics, **803-814**
 spinal cord stimulation, for angina pectoris and peripheral vascular disease, **795-802**
 outcomes assessment of therapeutic and cost benefit success rates, **815-821**
 intraoperative analgesia with intravenous ambulatory anesthesia, 276-278
 See also *Analgesia*.
- Pain mechanisms, 331-332
- Pain, postoperative, as patient outcome after ambulatory anesthesia, 407-408
- Paracervical block, during labor, 3-4
- Patent ductus arteriosus, in adult patients with congenital heart disease, 654, 659
- Patient education, prior to discharge after ambulatory surgery, 380-381
- Patient factors, incomplete epidural block for cesarean section due to, 46
- Patient outcomes, after ambulatory anesthesia, **403-415**
 economic outcomes, 411-412
 nontraditional, 409-411
 functional health status and quality of life, 409-410
 patient satisfaction, 410-411
 traditional, 404-409
 death, 404
 non-anesthetic-related, 404-405
 other perioperative complications, 408-409
 postoperative nausea and vomiting, 405-407
 postoperative pain, 407-408
- Patient perception, prior to discharge after ambulatory surgery, 380-381
- Patient satisfaction, with ambulatory anesthesia, 223-224
 as outcome measure, 410-411
- Patient selection, for ambulatory anesthesia, challenging patients, **243-263**
 elderly patients, 256-257
 infants, 257-258
 with aortic stenosis, 250-251
 with end-stage renal disease, 255-256
 with ischemic heart disease, 245-249
 with obstructive sleep apnea, 252-255
 for pediatric patients, 306-308
 history of, and developments in, 217-219
- Patient-controlled analgesia, during labor, epidural, 10-12
 inhalational, 2
 intravenous, 1-2
- Patient-controlled sedation, intravenous ambulatory anesthesia with, 281
- Pediatric ambulatory anesthesia, current issues in, **305-311**
 intraoperative considerations, 309-310
 patient selection, 306-308
 postoperative considerations, 310-311
 preoperative assessment, 308-309
 procedure selection, 305-306
- Pediatric patients, perioperative arrhythmias in, 580-584

- anesthetic factors, 581
- congenital heart disease and, 581-584
- Pegorgoten, for neuroprotection in cardiac surgery, 491-492
- Pelvic pain disorders, sacral neuromodulation for chronic, **783-793**
- Perception, patient, prior to discharge after ambulatory surgery, 380-381
- Perfluorochemical solutions, clinical potential during cardiac surgery, 554-555, 557-558
 - as transfusion alternative to red blood cells, 560-561
 - in cardiopulmonary bypass, 563
 - in myocardial contrast imaging, 561-562
 - in stroke, 564
 - protection against ischemia-reperfusion injury, 563-564
- Perioperative cardiac arrhythmias.
See Arrhythmias.
- Peripheral vascular disease, spinal cord stimulation for, **795-802**
- Pexelizumab, for neuroprotection in cardiac surgery, 492
- Pharmacy availability, for the cardiac catheterization laboratory, 645
- Phosphate depletion, intracellular high-energy, in perioperative myocardial ischemia-reperfusion injury, 466
- Phosphodiesterase inhibitors, for cardiopulmonary bypass-induced inflammation, 460
- Placenta, obstetric hemorrhage involving, abruptio placenta, 114-115
 - placenta accreta, 116
 - placenta previa, 115-116
 - retained placenta, 115
- Placenta accreta, obstetric hemorrhage due to, 116
- Placenta previa, obstetric hemorrhage due to, 115-116
- Planning, anesthesia, to limit and treat hemorrhage in obstetric patients, **127-144**
- Plasma inflammatory mediators, in systemic inflammatory response to cardiopulmonary bypass, 457-458
- Platelet activation, in systemic inflammatory response to cardiopulmonary bypass, 456
- Platelet function, for monitoring hemostasis after cardiopulmonary bypass, 519-520
- Platelets, pathophysiology of excessive bleeding after cardiopulmonary bypass, 529-530
- Podiatry, use of office-based anesthesia in, 438
- Point-of-care hemostasis monitors, after cardiopulmonary bypass, 520-523
- Postdural puncture headache, after regional analgesia during labor, **73-79**
- Postoperative nausea and vomiting, as patient outcome after ambulatory anesthesia, 405-407
 - management of, after ambulatory surgery, **347-365**
 - antiemetics, in clinical use, 349-355
 - with potential clinical use, 355-356
 - in pediatric patients, 310-311
 - nonpharmacologic techniques, 356-357
 - prediction of, 349
 - strategy for, 357-360
 - with inhalational anesthesia, 266-267
- Postoperative pain, as patient outcome after ambulatory anesthesia, 407-408
- Postoperative recovery and discharge, after ambulatory surgery, **367-386**
 - history of, and developments in, 216-217
- Postprocedural headache, with spinal anesthesia in outpatient surgery, 295
- Posttraumatic stress disorders, due to withholding of epidural analgesia during labor, 67-68
- Pre-eclampsia, **145-163**
 - anesthetic management, 156-160
 - epidural analgesia and, misconceptions about, 63-64
 - hemodynamic profile in, 146-153
 - treatment, 153-156
- Pre-emptive analgesia, 332
- Preconditioning, ischemic, in off-pump coronary artery bypass, 621
 - to diminish perioperative myocardial ischemia-reperfusion injury, 477-481
- Pregnancy-induced hypertension, obstetric emergencies due to, 120-121
- Pregnancy. *See also Labor and Obstetric anesthesia.*, gastrointestinal anatomy and physiology in, 90-91

- Preoperative assessment, for ambulatory surgery patients, **229-242**
 before day of surgery, 232-236
 history of, 215-216
 laboratory testing, 237-241
 patient and procedure selection, 232
 pediatric, 308-309
 personnel performing, 236-237
 philosophy of, 230-231
 risk classification, 231-232
- Preterm infants, ambulatory anesthesia in, 257-258
- Procaine, for spinal anesthesia in outpatient surgery, 297
- Professional recognition, of office-based anesthesia, 418-419
- Progress of labor, epidural analgesia and, misconceptions about, 60-63
- Prone position, use of laryngeal mask airway in, 321
- Propofol, for antiemetic use after ambulatory surgery, 354-355
 for neuroprotection in cardiac surgery, 488-489
 intravenous ambulatory anesthesia with, 275-276
- Prostodynia, sacral neuromodulation for, 787-792
 traditional medical and surgical therapies, 785-786
- Pruritus, as opioid side effect, 706
- Psychotherapy, impact of, on survival in cancer patients, 828
- Pulmonary aspiration, of gastric contents during labor, decline in maternal mortality due to, 88-90
- Pulmonary disease, in adults with congenital heart disease, 657
- Pulmonary embolism, during pregnancy, 166-170
- Pulmonary hypertension, in adults with congenital heart disease, 656-657
- Pulsatile pumps, in congestive heart failure, 628-629
- Q**
- Quality of life, as patient outcome after ambulatory anesthesia, 409-410

R

- Radiation safety, in the cardiac catheterization laboratory, 643-644
- Radiopaque media, in the cardiac catheterization laboratory, 646-647
- Re-entry, and perioperative arrhythmias, 572-574
- Reactive oxygen species, in perioperative myocardial ischemia-reperfusion injury, 466-467
 ischemic preconditioning before CABG with, 479
- Recombinant tissue plasminogen activator (r-tPA), for venous thromboembolism during pregnancy, 169
- Recovery, postoperative, after ambulatory surgery, **367-386**
 definitions of, 368-369
 discharge and, 372-376
 fast tracking, 369-371
 postoperative nausea and vomiting, 378-380
 postoperative pain, 376-378
 psychomotor tests of, 371
 with office-based anesthesia, 436-437
- Regional analgesia, techniques for, during labor, 3-12
 combined spinal-epidural analgesia, 5-10
 continuous spinal analgesia, 4-5
 epidural analgesia, patient-controlled, 10-12
 lumbar sympathetic block, 3
 paracervical block, 3-4
- Regional anesthesia, back pain after, **71-73**
 for outpatient surgery, **289-303**
 adverse events, 295-296
 continuous catheter techniques, 294
 epidural anesthesia, 298-299
 lower extremity blocks, 292-293
 neuraxial techniques, 294-295, 299-300
 spinal anesthesia, 296-298
 upper extremity blocks, 290-292
 headache after, **73-79**
 neurologic complications after, **79-82**
 cauda equina syndrome, 82
 etiology of, 79-80
 nerve injury from needle, 80-81
 transient neurologic syndrome, 81-82
 versus general, in cardiac catheterization laboratory, 644-645
- Regurgitation, intraoperative transesophageal echocardiography in, aortic, 600-602

- mitral, 591-593
tricuspid, 604-605
- Remacemide, for neuroprotection in cardiac surgery, 490-491
- Renal disease, end-stage, ambulatory anesthesia in patients with, 255-256
in adults with congenital heart disease, 657
- Renal dysfunction, in pre-eclampsia, 149
- Respiratory system changes, in pre-eclampsia, 151
- Resuscitation, in obstetric emergencies involving fetal compromise, in utero, 118
neonatal, 119
- Retained placenta, obstetric hemorrhage due to, 114-115
- Risk classification, in preoperative assessment for ambulatory surgery, 231-232
- Ropivacaine, compared with bupivacaine for obstetric analgesia, **19-38**
See also *Local anesthetics*.
- S**
-
- Sacral neuromodulation, for chronic pain conditions, **783-793**
- Safety issues, in office-based anesthesia, 426-428
- Saline injection, epidural, for postdural puncture headache, 77
- Salvage, of blood, intraoperative, for hemorrhage in obstetric patients, 134-135
- Satisfaction, patient, with ambulatory anesthesia, 223-224
as outcome measure, 410-411
recovery and discharge and, 380-381
surgeon, with ambulatory anesthesia, 223-224
- Scheduling, of cases, in an ambulatory surgical center, **387-401**
- Sciatic nerve block, distal, in outpatient surgery, 293
- Screening, preoperative. See *Preoperative assessment*.
- Sedation, as opioid side effect, 705
intravenous ambulatory anesthesia with, conscious, 279-280
patient-controlled, 281
- Serine protease inhibitors, for cardiopulmonary bypass-induced inflammation, 459-460
- Serotonin antagonists, for antiemetic use after ambulatory surgery, 353-354
- Sickle cell disease, effect of pain on survival in, 825
- Single ventricle, in adult patients with congenital heart disease, 656, 664-665
- Single-shot spinal anesthesia, after incomplete epidural block for cesarean section, 47-48
- Sleep apnea, obstructive, ambulatory anesthesia in patients with, 252-255
- Sodium nitroprusside, to control hypertension in pre-eclampsia, 154
- Somatostatin analogs, intraspinal analgesia with, 806
- Spastic disease states, with pain, botulinum toxin therapy for, 720
- Spinal analgesics, **803-814**
experimental agents, 809-811
intrathecal acetylcholinesterase inhibitors, 810
nitric oxide synthase inhibitors, 810
sustained release delivery of local anesthetics, 810-811
tricyclic antidepressants, 810
impact on survival, 829
intraspinal opioids, 804-806
fentanyl and sufentanil, 805
methadone and meperidine, 805
morphine, 805
side effects, 805-806
local anesthetics as adjuvant agents, 806
non-opioids, 806-809
alpha-2 adrenergic agents, 807-808
benzodiazepines, 809
calcium channel blockers, 808
cells in matrix, 809
long-acting local anesthetics, 809
N-methyl-D-aspartate (NMDA) antagonists, 808-809
somatostatin analogs, 806
spinal infusion research, 811-812
tolerance and safety, 811
treatment criteria, 803-804
- Spinal anesthesia, after incomplete epidural block for cesarean section, combined spinal-epidural technique, 49
continuous, 49
single-shot, 47-48

- combined epidural and, 5-10
- continuous, during labor, 4-5
- in outpatient surgery, 296-298
- Spinal blocks, differential, interventional pain management with, psychological aspects of, 774
- Spinal cord stimulation, for angina pectoris and peripheral vascular disease, **795-802**
 - complications, 800-801
 - mechanisms of action, 795-796
 - variables in, 796-797
 - outcomes assessment of therapeutic and cost benefit success rates, **815-821**
 - psychological aspects of interventional therapy with, 776-777
- Spinal-epidural analgesia, combined, during labor, 5-10
- Staffing issues, scheduling cases in an ambulatory surgery center, **387-401**
- Stenosis, aortic, ambulatory anesthesia in patients with, 250-251
 - intraoperative transesophageal echocardiography in, aortic, 599-600
 - mitral, 589-591
 - tricuspid, 604
- Steroids, for antiemetic use after ambulatory surgery, 354
- Streamlined Pharynx Airway Liner (SLIPA), 324
- Stroke, delayed/postoperative, and neuroprotection in cardiac surgery, 499-500
 - potential application of oxygen therapeutics in, 564
- Sufentanil. See also *Opioids*., intraspinal, 805
- Sumatriptan, for postdural puncture headache, 76
- Supplemental injections, epidural or caudal, after incomplete epidural block for cesarean section, 49-50
- Support services, anesthesia, for the cardiac catheterization laboratory, 645
- Supraclavicular block, in outpatient surgery, 291
- Supraventricular arrhythmias, perioperative, 574-577
 - acute management of, 574-576
 - chemical cardioversion of, 576-577
- Surgeons, general, use of office-based anesthesia by, 437
- satisfaction with ambulatory anesthesia, 223-224
- Survival, effect of pain on, **823-831**
 - impact of intrathecal opioid delivery, 829
 - impact of pain management therapies-829, 828
 - impact of psychotherapy on, 827-828
 - in cancer, 825-827
 - in coronary artery disease, 823-825
 - in sickle cell disease, 825
 - interaction of pain and mood, 827
- Survival rates, long-term, in cardiac surgery, gender differences in, 683-684
- Sustained-release delivery, intrathecal, of local anesthetics, 810-811
- Systemic inflammatory response, to cardiopulmonary bypass, **453-464**
 - anti-inflammatory therapeutic strategies, 458-462
 - mechanical, 461-462
 - pharmacologic, 459-461
 - cellular response, 455-456
 - endothelial activation, 456
 - leukocyte activation, 455-456
 - platelet activation, 456
 - contact activation, 453-455
 - complement system, 453-455
 - factor XII/kallikrein system, 455
 - plasma inflammatory mediators, 457-458
 - arachidonic acid metabolites, 458
 - cytokines and chemokines, 457
 - endothelin-1, 458
 - endotoxemia, 458
 - nitric oxide, 458
- Systemic inflammatory response syndrome (SIRS), in perioperative myocardial ischemia-reperfusion injury, 474

T

- Target-controlled infusion, intravenous ambulatory anesthesia with, 279
- Tetralogy of Fallot, in adult patients with congenital heart disease, 655, 662-663
- Theophylline, for postdural puncture headache, 76

- Thiopental, for neuroprotection in cardiac surgery, 487-488
intravenous ambulatory anesthesia with, 274
- Thoracic epidural neuroplasty, 762-763
- Thrombin inhibitors, monitoring of, for cardiopulmonary bypass, 517
- Thrombin time, high dose, monitoring of, for cardiopulmonary bypass, 516
- Thrombocytopenia, obstetric anesthesia for patients with, **99-109**
low-molecular-weight heparin and, 105-107
practical recommendations, 104-105
- Thrombolysis, for venous thromboembolism during pregnancy, 169-170
- Tissue plasminogen activator, recombinant (r-tPA), for venous thromboembolism during pregnancy, 169
- Total intravenous anesthesia (TIVA), 278-279
- Toxicity, organ, as opioid side effect, 706
- Tranexamic acid, use of during routine cardiopulmonary bypass, 533-536
- Transcutaneous electrical nerve stimulation (TENS), for pain management in ambulatory setting, 342
- Transesophageal echocardiography, intraoperative, in valvular heart surgery, **587-612**
aortic valve procedures, 596-603
detection of intracardiac air, 607-608
mitral valve procedures, 588-596
tricuspid valve procedures, 604-607
- Transfusions, oxygen therapeutics as alternative to red blood cells for, 559-561
- Transfusions, blood, risks of, in obstetric cases, 128
future alternatives to, 138-139
- Transient neurologic syndrome, after regional anesthesia, 81-82
- Transposition of great vessels, in adult patients with congenital heart disease, 655, 660-661
congenitally corrected, 655, 661
- Tricuspid regurgitation, intraoperative transesophageal echocardiography in, 604-605
- Tricuspid stenosis, intraoperative transesophageal echocardiography in, 604
- Tricuspid valve procedures, transesophageal echocardiography in, 604-607
after cardiopulmonary bypass, 606-607
before cardiopulmonary bypass, 604-606
- Tricyclic antidepressants, intraspinal analgesia with, 810
- Trimethaphan, to control hypertension in pre-eclampsia, 154
- Truncus arteriosus, in adult patients with congenital heart disease, 655, 663
-
- U**
- Upper extremity blocks, for regional anesthesia in outpatient surgery, 290-292
- Upper motor neuron disease syndromes, with pain, botulinum toxin therapy for, 720-721
- Urology, use of office-based anesthesia in, 438
- Uterine atony, obstetric hemorrhage due to, 112-114
- Uterine inversion, obstetric hemorrhage due to, 117
- Uterine rupture, obstetric hemorrhage due to, 116-117
-
- V**
- Valvular anomalies, in adult patients with congenital heart disease, 660
- Valvular heart surgery, transesophageal echocardiography during, **587-612**
aortic valve procedures, 596-603
detection of intracardiac air, 607-608
mitral valve procedures, 588-596
tricuspid valve procedures, 604-607
- Vascular disease, peripheral, spinal cord stimulation for, **795-802**
- Vasodilators, direct-acting, to control hypertension in pre-eclampsia, 154
- Venous air embolism, during pregnancy, 170-173
- Venous thromboembolism, during pregnancy, 166-170
- Ventricle, single, in adult patients with congenital heart disease, 656, 664-665
- Ventricular arrhythmias, perioperative, 577-580

- drug selection for acute management of, 579-580
 - nonsustained, 577-578
 - sustained, 578-579
- Ventricular assist devices, left, **625-638**
- anesthetic management for insertion of, 633-637
 - historical evolution and indications, 626-627
 - physiology and preoperative considerations, 630-632
 - surgical approach, 633
 - types of, 627-630
 - counterpulsation, 627
 - extracorporeal kinetic displacement, 627-628
 - nonpulsatile pumps, 629-630
 - pulsatile pumps, 628-629
- Ventricular dysfunction, in adults with congenital heart disease, 657
- Ventricular septal defect, in adult patients with congenital heart disease, 654, 658-659
- Voiding difficulty, with neuraxial anesthesia in outpatient surgery, 296
- Vomiting, as opioid side effect, 703
- See also *Nausea and vomiting, postoperative.*
- Vulvodynia, sacral neuromodulation for, 787-792
- traditional medical and surgical therapies, 785-786

W

- Women, gender differences in cardiac surgery, **675-689**
- operative factors and mortality, 682-683
 - pharmacology and estrogen replacement therapy, 679-682
 - postoperative long-term follow-up, 683-684
 - preoperative risk profile and mortality, 675-678
 - role of bias, 678-679
- Writer's cramp, botulinum toxin therapy for, 719-720